

**Table 6D**  
**2003 Pediatric Nutrition Surveillance** (1)  
**National**  
**Comparison of Growth and Anemia Indicators by Contributor**  
**Children Aged < 5 Years** (2)

Page: 1  
Run Date: 7 / 9 / 04

<b>Comparison of Growth and Anemia Indicators by Contributor</b> (3,4)											
<b>Contributor</b>	<b>Birthweight (5)</b>			<b>Height and Weight</b>						<b>Anemia Low Hb/Hct (8)</b>	
	<b>Low</b>		<b>High</b>	<b>Short Stature (6)</b>			<b>Under- weight (7)</b>	<b>Over- weight (7)</b>	<b>≥ 2 Yrs Over- weight (7)</b>		
	<b>Number</b>	<b>% &lt;2500 g (Rank)</b>	<b>% &gt;4000 g (Rank)</b>	<b>Number</b>	<b>% &lt;5th (Rank)</b>	<b>% &lt;5th (Rank)</b>	<b>% ≥95th (Rank)</b>	<b>Number</b>	<b>% 85th- &lt;95th (Rank)</b>	<b>% ≥95th (Rank)</b>	<b>Number</b> <b>% (Rank)</b>
Alabama	0	* *	* *	19,444	6.9(26)	4.9(30)	14.5(34)	8,235	16.4(28)	14.7(32)	9,537      22.7 (44)
Arkansas	20,281	9.5 (29)	6.7 (13)	72,110	7.3(35)	5.1(32)	10.8 (5)	31,625	14.5(11)	12.2(11)	49,931      11.3 (20)
AZ I/Tribal Council	2,711	6.3 (2)	9.0 (33)	11,001	6.8(24)	2.6 (4)	20.9(42)	5,037	20.9(41)	20.9(43)	7,706      12.3 (25)
California	166,630	7.1 (6)	8.6 (31)	871,685	4.9 (9)	5.4(35)	16.2(36)	344,384	16.2(26)	17.6(40)	500,508      13.7 (31)
Cheyenne River-SD	138	9.4 (27)	8.7 (32)	745	4.0 (3)	1.5 (1)	25.1(44)	388	20.9(41)	17.5(39)	425      11.3 (20)
Chickasaw Nation-OK	924	8.7 (22)	9.1 (35)	3,313	6.9(26)	3.9(16)	11.5 (8)	1,478	14.3 (8)	12.0 (9)	2,187      17.9 (42)
Colorado	32,258	10.2 (35)	4.9 (3)	118,648	7.9(39)	4.2(19)	8.6 (3)	50,773	13.6 (5)	9.4 (2)	51,116      7.8 (5)
D.C.	4,200	11.9 (41)	5.8 (5)	13,976	7.2(33)	5.3(33)	11.8(12)	5,926	14.2 (7)	13.3(21)	10,199      21.0 (43)
Florida	111,536	9.6 (30)	6.6 (11)	383,505	4.6 (7)	5.6(37)	12.3(18)	155,483	14.8(15)	13.4(24)	255,639      14.8 (35)
Georgia	73,525	10.0 (33)	6.4 (10)	248,553	7.2(33)	4.3(21)	13.3(28)	92,728	14.4 (9)	12.4(12)	150,648      13.4 (29)
Hawaii	8,099	10.1 (34)	7.4 (23)	36,370	7.1(31)	3.4(13)	12.7(21)	15,602	12.9 (2)	10.1 (4)	16,465      10.4 (15)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6D**  
**2003 Pediatric Nutrition Surveillance** (1)  
**National**  
**Comparison of Growth and Anemia Indicators by Contributor**  
**Children Aged < 5 Years** (2)

Page: 2  
Run Date: 7 / 9 / 04

Comparison of Growth and Anemia Indicators by Contributor (3,4)													
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Idaho	9,603	6.9 (4)	7.5 (25)	36,647	6.7(23)	4.4(24)	9.8 (4)	16,341	14.7 (13)	11.1 (6)		24,904	12.8 (26)
Illinois	53,941	9.1 (24)	7.2 (18)	186,123	7.1(31)	4.4(24)	13.1(25)	70,617	15.3(17)	14.0(30)		106,677	10.9 (17)
Indiana	38,580	9.0 (23)	7.3 (22)	126,651	6.5(22)	4.2(19)	11.5 (8)	51,953	16.2(26)	13.7(29)		77,646	15.3 (37)
Iowa	6,776	7.1 (6)	9.0 (33)	45,133	5.9(15)	2.5 (3)	13.9(32)	21,892	17.7(35)	13.6(26)		23,685	10.1 (13)
Kansas	16,098	8.4 (20)	7.0 (16)	63,614	6.9(26)	3.0 (9)	12.5(19)	27,076	16.0(23)	12.6(15)		43,683	10.9 (17)
Kentucky	26,648	9.6 (30)	7.4 (23)	121,974	7.4(36)	4.0(17)	17.7(39)	60,984	17.8(37)	17.2(37)		79,078	11.5 (23)
Louisiana	33,877	11.9 (41)	5.5 (4)	111,763	9.3(43)	5.4(35)	13.0(24)	44,036	15.0(16)	13.3(21)		78,656	14.9 (36)
Maine	5,766	7.8 (12)	11.4 (42)	21,872	7.6(38)	6.3(38)	13.6(31)	9,876	16.8(31)	16.0(35)		13,412	12.9 (27)
Michigan	50,792	9.4 (27)	8.3 (28)	216,362	6.9(26)	4.3(21)	12.7(21)	93,962	16.1(24)	12.9(16)		134,250	13.1 (28)
Minnesota	27,877	7.4 (9)	10.0 (39)	93,609	4.2 (4)	3.3(10)	13.5(30)	40,161	15.9(20)	13.2(19)		74,648	9.8 (11)
Missouri	39,423	9.1 (24)	6.9 (15)	131,869	6.9(26)	4.6(28)	12.0(13)	56,346	16.4(28)	13.3(21)		86,726	16.2 (39)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6D**  
**2003 Pediatric Nutrition Surveillance** (1)  
**National**  
**Comparison of Growth and Anemia Indicators by Contributor**  
**Children Aged < 5 Years** (2)

Page: 3  
Run Date: 7 / 9 / 04

Comparison of Growth and Anemia Indicators by Contributor (3,4)													
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Montana	3,927	7.9 (14)	8.4 (29)	21,338	4.9 (9)	2.9 (8)	12.0(13)	10,178	15.7 (19)	11.0 (5)		12,821	8.8 (9)
Navajo Nation	3,146	6.7 (3)	7.0 (16)	14,937	5.3(14)	2.7 (6)	13.1(25)	7,616	18.5(39)	14.4(31)		11,700	8.3 (7)
Nebraska	10,964	8.0 (15)	7.7 (26)	41,671	6.1(17)	4.4(24)	13.2(27)	17,242	16.1 (24)	13.4(24)		25,792	13.6 (30)
Nevada	10,811	8.4 (20)	7.2 (18)	38,731	8.1(40)	2.8 (7)	14.1(33)	14,595	15.9(20)	13.6(26)		27,221	10.4 (15)
New Hampshire	4,038	7.8 (12)	10.8 (41)	15,865	6.8(24)	3.3(10)	16.3(37)	7,227	19.4(40)	15.6(34)		10,490	13.8 (32)
New Jersey	40,257	9.1 (24)	7.2 (18)	142,092	6.4(21)	5.0(31)	16.9(38)	56,774	17.0(32)	17.9(41)		102,762	16.9 (40)
New York	70,739	7.6 (11)	8.0 (27)	409,105	4.4 (6)	7.2(42)	13.4(29)	178,642	16.5(30)	16.8(36)		274,354	14.1 (33)
North Dakota	3,182	7.0 (5)	10.7 (40)	13,495	4.2 (4)	3.5(14)	12.1(15)	6,106	15.6(18)	11.4 (7)		8,058	7.7 (4)
Ohio	56,684	10.2 (35)	6.7 (13)	207,992	6.1(17)	4.1(18)	12.5(19)	89,824	14.7(13)	11.6 (8)		129,607	14.2 (34)
Oregon	10,179	5.6 (1)	19.7 (43)	52,380	6.0(16)	2.6 (4)	15.8(35)	29,875	17.2(34)	14.7(32)		42,507	11.3 (20)
Pennsylvania	50,496	10.3 (37)	7.2 (18)	226,988	6.1(17)	6.5(40)	11.7(11)	100,053	14.4 (9)	12.4(12)		162,362	15.3 (37)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6D**  
**2003 Pediatric Nutrition Surveillance** (1)  
**National**  
**Comparison of Growth and Anemia Indicators by Contributor**  
**Children Aged < 5 Years** (2)

Page: 4  
Run Date: 7 / 9 / 04

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Puerto Rico	42,788	11.6 (40)	2.9 (1)	207,534	9.0(42)	8.3(44)	20.1(41)	102,624	18.3(38)	24.0(44)	168,221	8.3 (7)
Rosebud Sioux-SD	231	7.4 (9)	9.5 (37)	1,325	3.2 (2)	1.8 (2)	18.0(40)	641	21.5(43)	17.3(38)	1,062	17.8 (41)
South Carolina	21,040	12.5 (43)	6.1 (6)	83,775	13.2(44)	7.9(43)	11.0 (7)	32,239	13.5 (4)	12.4(12)	57,388	10.9 (17)
South Dakota	4,968	8.0 (15)	9.4 (36)	19,578	6.1(17)	3.5(14)	12.7(21)	8,423	17.7(35)	13.6(26)	14,018	7.6 (3)
Standing Rock-ND	178	7.3 (8)	6.2 (8)	840	1.9 (1)	6.4(39)	22.3(43)	422	22.0(44)	20.1(42)	652	8.0 (6)
Tennessee	44,412	10.4 (38)	6.1 (6)	146,390	5.2(12)	6.8(41)	10.9 (6)	60,086	13.8 (6)	12.0 (9)	95,067	7.1 (2)
Utah	17,798	8.0 (15)	6.3 (9)	69,368	7.5(37)	4.3(21)	7.6 (1)	31,099	12.6 (1)	8.6 (1)	47,406	9.9 (12)
Vermont	2,756	8.2 (18)	9.9 (38)	16,293	5.1(11)	3.3(10)	12.2(16)	8,504	17.1(33)	13.1(18)	10,311	9.0 (10)
West Virginia	13,002	9.7 (32)	6.6 (11)	51,809	4.8 (8)	5.3(33)	11.5 (8)	22,079	14.5(11)	13.2(19)	34,010	6.6 (1)
Wisconsin	28,471	8.3 (19)	8.5 (30)	109,384	5.2(12)	4.4(24)	12.2(16)	50,284	15.9(20)	13.0(17)	75,448	12.2 (24)
Wyoming	3,116	10.7 (39)	4.2 (2)	12,400	8.7(41)	4.8(29)	7.9 (2)	5,269	12.9 (2)	9.5 (3)	9,058	10.3 (14)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6D**  
**2003 Pediatric Nutrition Surveillance** (1)  
**National**  
**Comparison of Growth and Anemia Indicators by Contributor**  
**Children Aged < 5 Years** (2)

Page: 5  
Run Date: 7 / 9 / 04

Comparison of Growth and Anemia Indicators by Contributor (3,4)													
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Nation	1,172,866	9.1	7.3	4,838,257	6.2	5.2	13.6	2,044,705	15.7	14.7		3,118,041	12.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.